

Topics covered in this tutorial

- Conditional PMFs/PDFs and expectations
- Gaussian conditioning and independence
- Gaussian vectors, covariance and correlation matrices
- Affine transformations and cross-covariance

Complete Questions 1–5 before the tutorial test (normally around 16:00–16:15). Questions 6–8 are retained at the end for further practice.

Onderwerpe wat in hierdie tutoriaal gedek word

- Voorwaardelike WMF's/WDF's en verwagtings
- Gaussiese kondisionering en onafhanklikheid
- Gaussiese vektore, kovariansie- en korrelasiematrikse
- Affiene transformasies en kruiskovariansie

Voltooi Vrae 1–5 voor die tutoriaaltoets (gewoonlik omstreeks 16:00–16:15). Vrae 6–8 word aan die einde vir verdere oefening behou.

1. Let $\mathbf{X} = [X_1 \ X_2]'$ be a discrete random vector. Each component takes values in $\{1, 2, 3\}$, so its support is

$$S_{\mathbf{X}} = \{\mathbf{x} = [x_1 \ x_2]' : x_1, x_2 \in \{1, 2, 3\}\}.$$

The joint probability mass function is

$$P_{\mathbf{X}}(\mathbf{x}) = P_{\mathbf{X}}([x_1 \ x_2]') = P_{X_1, X_2}(x_1, x_2) = \begin{cases} \frac{x_1 + x_2}{36}, & \mathbf{x} \in S_{\mathbf{X}}, \\ 0, & \mathbf{x} \notin S_{\mathbf{X}}. \end{cases}$$

- (a) Find the complete PMF of $S = X_1 + X_2$. Hence find $P(S = 3)$.
 (b) Find the marginal PMF $P_{X_2}(j)$ and the conditional PMF $P_{X_1|X_2}(i | j)$ for $j \in \{1, 2, 3\}$.
 (c) Calculate $E[X_1 | X_2 = j]$. Use this conditional expected value to argue whether X_1 and X_2 are independent.
 (d) Use the PMF to calculate $\boldsymbol{\mu}_{\mathbf{X}}$, $\mathbf{R}_{\mathbf{X}} = E[\mathbf{X}\mathbf{X}']$ and $\mathbf{C}_{\mathbf{X}}$. Use symmetry to reduce the calculations.

2. The continuous random variables X and Y have joint PDF

$$f_{X,Y}(x, y) = \begin{cases} \frac{1}{2}, & -1 \leq x \leq y \leq 1, \\ 0, & \text{otherwise / andersins.} \end{cases}$$

- (a) Find the marginal PDF $f_Y(y)$.
 (b) Find $f_{X|Y}(x | y)$ and state the values of y for which the conditional-density formula is valid.
 (c) Calculate $E[X | Y = y]$ and $E[Y - X | Y = y]$.
 (d) Use iterated expectation to calculate $E[X]$.

- Laat $\mathbf{X} = [X_1 \ X_2]'$ 'n diskrete toevalsvektor wees. Elke komponent neem waardes in $\{1, 2, 3\}$ aan, dus is die steunversameling

Die gesamentlike waarskynlikheidsmassafunksie is

- (a) Vind die volledige WMF van $S = X_1 + X_2$. Bepaal dus $P(S = 3)$.
 (b) Vind die marginale WMF $P_{X_2}(j)$ en die voorwaardelike WMF $P_{X_1|X_2}(i | j)$ vir $j \in \{1, 2, 3\}$.
 (c) Bereken $E[X_1 | X_2 = j]$. Gebruik hierdie voorwaardelike verwagtingswaarde om te motiveer of X_1 en X_2 onafhanklik is.
 (d) Gebruik die WMF om $\boldsymbol{\mu}_{\mathbf{X}}$, $\mathbf{R}_{\mathbf{X}} = E[\mathbf{X}\mathbf{X}']$ en $\mathbf{C}_{\mathbf{X}}$ te bereken. Gebruik simmetrie om die berekeninge te verminder.

Die kontinue toevalsveranderlikes X en Y het gesamentlike WDF

- (a) Vind die marginale WDF $f_Y(y)$.
 (b) Vind $f_{X|Y}(x | y)$ en meld die waardes van y waarvoor die voorwaardelike-digtheidsformule geldig is.
 (c) Bereken $E[X | Y = y]$ en $E[Y - X | Y = y]$.
 (d) Gebruik herhaalde verwagting om $E[X]$ te bereken.

3. A temperature-monitoring system records temperature T (in °C) and a sensor voltage W (in V). They are jointly Gaussian, with the means and *standard deviations* below. An alarm occurs on the event $I = \{T > 38, W > 10\}$.

$$\mu_T = 37, \quad \sigma_T = 1,$$

(a) First assume $\rho_{T,W} = 0$. Explain why the individual probabilities can be multiplied as follows, then calculate $P(I)$.

$$P(I) = P(T > 38, W > 10) = P(T > 38) P(W > 10).$$

(b) Now use $\rho_{T,W} = 1/\sqrt{2}$ for the remaining parts. Find the conditional mean, variance and PDF of W given $T = t$.

(c) Express $P(I | T = t)$ in terms of Φ (the standard Gaussian CDF) for *all* real t . No numerical evaluation is required.

(d) Compare the conditional variance with $\text{Var}[W]$. What does this say about the uncertainty in W once T is known?

4. $\mathbf{X}$ is a 3-dimensional Gaussian random vector with mean vector $\boldsymbol{\mu}_X = [4 \ 8 \ 6]'$ and covariance matrix

$$\mathbf{C}_X = \begin{bmatrix} 4 & -2 & 1 \\ -2 & 4 & -2 \\ 1 & -2 & 4 \end{bmatrix}.$$

(a) Calculate the correlation matrix $\mathbf{R}_X = \mathbf{E}[\mathbf{X}\mathbf{X}']$.

(b) Find the joint PDF $f_{X_1, X_2}(x_1, x_2)$.

(c) Calculate $P(X_1 > 8)$. Use $\Phi(2) = 0.9772$.

(d) Are X_1, X_2, X_3 independent? Justify using the covariance matrix and the Gaussian assumption.

5. Given the Gaussian random vector $\mathbf{X}$ in Question 4, define $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 1/2 & 2/3 \\ 1 & -1/2 & 2/3 \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} -4 \\ -4 \end{bmatrix}$$

Calculate:

- (a) $\boldsymbol{\mu}_Y$ (b) $\mathbf{C}_Y$ (c) $\mathbf{R}_Y$
(d) $P(-1 \leq Y_2 \leq 1)$ (e) $\mathbf{C}_{XY}$ (f) $\mathbf{R}_{XY}$
(g) ρ_{X_1, Y_2}

For (d), briefly justify why Y_2 is Gaussian. An answer in terms of Φ is sufficient.

(h) Can $\mathbf{C}_X$ be recovered here using $\mathbf{A}^{-1}\mathbf{C}_Y(\mathbf{A}^{-1})'$? Explain using the dimensions of $\mathbf{A}$.

'n Temperatuurmoniteringstelsel meet temperatuur T (in °C) en 'n sensorspanning W (in V). Hulle is gesamentlik Gaussies, met die gemiddeldes en *standaardafwykings* hieronder. 'n Alarm vind plaas tydens die gebeurtenis $I = \{T > 38, W > 10\}$.

$$\mu_W = 7, \quad \sigma_W = 2.$$

(a) Aanvaar eers $\rho_{T,W} = 0$. Verduidelik waarom die afsonderlike waarskynlikhede soos volg vermenigvuldig kan word, en bereken dan $P(I)$.

(b) Gebruik nou $\rho_{T,W} = 1/\sqrt{2}$ vir die oorblywende dele. Vind die voorwaardelike gemiddelde, variansie en WDF van W gegee $T = t$.

(c) Druk $P(I | T = t)$ in terme van Φ (die standaard-Gaussiese KVF) uit vir *alle* reële t . Geen numeriese berekening word vereis nie.

(d) Vergelyk die voorwaardelike variansie met $\text{Var}[W]$. Wat sê dit oor die onsekerheid in W wanneer T bekend is?

$\mathbf{X}$ is 'n 3-dimensionele Gaussiese toevalsvektor met gemiddeldevektor $\boldsymbol{\mu}_X = [4 \ 8 \ 6]'$ en kovariansiematriks

(a) Bereken die korrelasiematriks $\mathbf{R}_X = \mathbf{E}[\mathbf{X}\mathbf{X}']$.

(b) Vind die gesamentlike WDF $f_{X_1, X_2}(x_1, x_2)$.

(c) Bereken $P(X_1 > 8)$. Gebruik $\Phi(2) = 0.9772$.

(d) Is X_1, X_2, X_3 onafhanklik? Motiveer met die kovariansiematriks en die Gaussiese aanname.

Gegewe die Gaussiese toevalsvektor $\mathbf{X}$ in Vraag 4, definieer $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$, waar

$$\mathbf{A} = \begin{bmatrix} 1 & 1/2 & 2/3 \\ 1 & -1/2 & 2/3 \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} -4 \\ -4 \end{bmatrix}$$

Bereken:

- (a) $\boldsymbol{\mu}_Y$ (b) $\mathbf{C}_Y$ (c) $\mathbf{R}_Y$
(d) $P(-1 \leq Y_2 \leq 1)$ (e) $\mathbf{C}_{XY}$ (f) $\mathbf{R}_{XY}$
(g) ρ_{X_1, Y_2}

Motiveer vir (d) kortliks waarom Y_2 Gaussies is. 'n Antwoord in terme van Φ is voldoende.

(h) Kan $\mathbf{C}_X$ hier herwin word met $\mathbf{A}^{-1}\mathbf{C}_Y(\mathbf{A}^{-1})'$? Verduidelik aan die hand van $\mathbf{A}$ se afmetings.

Further practice

These questions are retained as further practice alongside the main tutorial (Questions 1–5).

6. The random vector $\mathbf{X} = [X_1 \ X_2 \ X_3]'$ has joint PDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} (x_1 + 2x_2 + 3x_3)/3, & 0 \leq x_1, x_2, x_3 \leq 1, \\ 0, & \text{otherwise / andersins.} \end{cases}$$

(a) Find $f_{X_1, X_2}(x_1, x_2)$ and $f_{X_3}(x_3)$. Is the subvector $[X_1 \ X_2]'$ independent of X_3 ? Use factorisation to justify your answer.

(b) Derive $F_{\mathbf{X}}(\mathbf{x})$ for $0 \leq x_1, x_2, x_3 \leq 1$. Calculate $F_{\mathbf{X}}(0.3, 0.5, 1)$.

(c) Calculate $E[X_1 X_2 X_3]$ directly from the joint PDF.

7. Three tasks start simultaneously. Their completion times T_1, T_2, T_3 are independent and identically distributed, each uniform on $[0, 2]$ seconds. Let $M = \max(T_1, T_2, T_3)$ and $L = \min(T_1, T_2, T_3)$.

(a) Derive the complete CDF and PDF of M , starting from the event that all three tasks finish by time t .

(b) Derive the complete CDF and PDF of L .

(c) Find the probability that all tasks finish within 1.5 s, and the probability that at least one finishes within 0.5 s.

(d) Use independence to find $E[T_1 T_2 T_3]$.

8. X_1 and X_2 are independent, each uniform on $[0, 1]$. Define $Y_1 = X_1 + X_2 + 1$ and $Y_2 = X_1 - X_2$.

(a) Write $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$. Find $\mathbf{A}^{-1}$ and $\det(\mathbf{A})$, and express x_1, x_2 in terms of y_1, y_2 .

(b) Sketch the support of $\mathbf{Y}$ and derive its joint PDF using the vector-transformation formula.

(c) Given $\mathbf{C}_{\mathbf{X}} = \frac{1}{12}\mathbf{I}_2$, where $\mathbf{I}_2$ is the 2×2 identity matrix, calculate $\mathbf{C}_{\mathbf{Y}}$. Recover $\mathbf{C}_{\mathbf{X}}$ using $\mathbf{A}^{-1}$. Are Y_1 and Y_2 independent? Justify using the support in (b).

(d) Consider the *separate* transformation $V_i = 2X_i + 1$, $i = 1, 2$. Express $F_{\mathbf{V}}(v_1, v_2)$ in terms of $F_{\mathbf{X}}$, then give the joint PDF and support of $\mathbf{V}$.

Verdere oefening

Hierdie vrae word as verdere oefening saam met die hooftutoriaal (Vrae 1–5) behou.

Die toevalsvektor $\mathbf{X} = [X_1 \ X_2 \ X_3]'$ het gesamentlike WDF

$$f_{\mathbf{X}}(\mathbf{x}) = \begin{cases} (x_1 + 2x_2 + 3x_3)/3, & 0 \leq x_1, x_2, x_3 \leq 1, \\ 0, & \text{otherwise / andersins.} \end{cases}$$

(a) Vind $f_{X_1, X_2}(x_1, x_2)$ en $f_{X_3}(x_3)$. Is die subvektor $[X_1 \ X_2]'$ onafhanklik van X_3 ? Gebruik faktoriserings om jou antwoord te motiveer.

(b) Lei $F_{\mathbf{X}}(\mathbf{x})$ af vir $0 \leq x_1, x_2, x_3 \leq 1$. Bereken $F_{\mathbf{X}}(0.3, 0.5, 1)$.

(c) Bereken $E[X_1 X_2 X_3]$ direk uit die gesamentlike WDF.

Drie take begin gelyktydig. Hul voltooiingstye T_1, T_2, T_3 is onafhanklik en identies verdeel, elk uniform op $[0, 2]$ sekondes. Laat $M = \max(T_1, T_2, T_3)$ en $L = \min(T_1, T_2, T_3)$ wees.

(a) Lei die volledige KVF en WDF van M af, vanaf die gebeurtenis dat al drie take teen tyd t klaar is.

(b) Lei die volledige KVF en WDF van L af.

(c) Vind die waarskynlikheid dat alle take binne 1.5 s klaar is, en die waarskynlikheid dat minstens een binne 0.5 s klaar is.

(d) Gebruik onafhanklikheid om $E[T_1 T_2 T_3]$ te vind.

X_1 en X_2 is onafhanklik, elk uniform op $[0, 1]$. Definieer $Y_1 = X_1 + X_2 + 1$ en $Y_2 = X_1 - X_2$.

(a) Skryf $\mathbf{Y} = \mathbf{A}\mathbf{X} + \mathbf{b}$. Vind $\mathbf{A}^{-1}$ en $\det(\mathbf{A})$, en druk x_1, x_2 in terme van y_1, y_2 uit.

(b) Skets die steunversameling van $\mathbf{Y}$ en lei sy gesamentlike WDF af met die vektortransformasieformule.

(c) Gegee $\mathbf{C}_{\mathbf{X}} = \frac{1}{12}\mathbf{I}_2$, waar $\mathbf{I}_2$ die 2×2 -identiteitsmatriks is, bereken $\mathbf{C}_{\mathbf{Y}}$. Herwin $\mathbf{C}_{\mathbf{X}}$ met $\mathbf{A}^{-1}$. Is Y_1 en Y_2 onafhanklik? Motiveer met die steunversameling in (b).

(d) Beskou die *afsonderlike* transformasie $V_i = 2X_i + 1$, $i = 1, 2$. Druk $F_{\mathbf{V}}(v_1, v_2)$ in terme van $F_{\mathbf{X}}$ uit, en gee dan die gesamentlike WDF en steunversameling van $\mathbf{V}$.